



ISSN 0972-060X

**Composition and Antioxidant Activities of the Essential Oil of Cinnamon
(*Cinnamomum zeylanicum* Blume) Leaves from Sri Lanka**

Erich Schmidt*

Kurt Kitzing Co., Hinterm Alten Schloss 21, D-86757 Wallerstein, Germany

Leopold Jirovetz, Gerhard Buchbauer and Gernot A. Eller

University of Vienna, Department of Clinical Pharmacy and Diagnostics,
Althanstrasse 14, A-1090 Vienna, Austria

Ivanka Stoilova and Albert Krastanov

University of Food Technologies, Department of Biotechnology, 26 Maritza Boulevard,
4002 Plovdiv, Bulgaria

Albena Stoyanova

University of Food Technologies, Department of Essential Oils,
26 Maritza Boulevard, 4002 Plovdiv, Bulgaria

Margit Geissler

Shimadzu-Germany, Department of GC and GC-MS, Albert-Hahn-Strasse 6-10,
D-47269 Duisburg, Germany

Received 20 May 2006; accepted 7 July 2006

Abstract: The composition of the volatiles from leaves of *Cinnamomum zeylanicum* Blume from Sri Lanka was studied by GC-FID and GC-MS. The basic component of the oil was found to be eugenol (74.9%), followed by β -caryophyllene (4.1%), benzyl benzoate (3.0%), linalool (2.5%), eugenyl acetate (2.1%) and cinnamyl acetate (1.8%). The essential leaf oil from cinnamon demonstrated scavenger activity against the DPPH radical at concentrations which are lower than the concentrations of eugenol, butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA). This essential cinnamon oil showed also a significant inhibitory effect on hydroxyl radicals and acted as an iron chelator. Cinnamon leaf oil efficiently inhibited the formation of conjugated dienes and the generation of secondary products from lipid peroxidation at a concentration equivalent to that of the standard BHT.

Key words: *Cinnamomum zeylanicum* Blume, Lauraceae, essential leaf oil, composition of volatiles, antioxidant activities

*Corresponding author: (Erich Schmidt)

E-mail: <erich.schmidt@kurtkitzing.de>

Introduction: *Cinnamomum zeylanicum* Blume syn. *C. verum* J. Presl, *Laurus cinnamomum* L. is an evergreen tropical tree, belonging to the Lauraceae family. Cinnamon barks and leaves are widely used as spice and flavoring agent in foods³⁴ and for various applications in medicine⁴⁰. From different parts of the plant (bark, roots and leaves) essential oils with variations in the composition by especially geographic and technical reasons can be obtained. The essential oil from *C. zeylanicum* barks is rich in *trans*-cinnamaldehyde with antimicrobial effects against animal and plant pathogens, food poisoning and spoilage bacteria and fungi^{6,26,29}. Until now more than 300 volatiles were found as constituents of essential oils of cinnamon⁴. The essential oil derived from cinnamon leaves is rich in eugenol, that from the roots in camphor and that from the barks shows a high amount of sesquiterpenes (α -bergamotene and α -copaene)¹⁹. It has been established that the oils and extracts from cinnamon possess a distinct antioxidant activity, which is especially attributed to the presence of phenolic and polyphenolic substances^{8,9,20,21,25,27,31,35,39}. Free radicals are generated in consequence of the metabolism of normal or pathological cells.

The electron acceptor molecule oxygen counteracts the free radicals transforming them into a reactive oxygen species (ROS). The primary derivatives of oxygen, such as superoxide anion radicals ($O_2^{\cdot-}$), hydroxyl radicals (OH^{\cdot}), hydrogen peroxide (H_2O_2) and the singlet oxygen (1O_2), play an important role in mediating ROS-related effects. Short-living reactive species generated *in situ* can react with non-radicals and produce chain reactions¹⁴. In the last years there has been a constantly rising interest in the antioxidative constituents of various medicinal plants, due to their potential in controlling the levels of free radicals and the process of lipid peroxidation^{32,38}.

Now, research is focused on the extraction of natural antioxidants (e.g. also from cinnamon)³³ that would be less toxic and more effective than the commonly applied synthetic antioxidants to fight the so-called "oxidative stress", which is a highly pathological situation⁵. In living organisms the protection against free radicals and reactive oxygen species is realized by antioxidative enzymes¹¹, reduced glutathione, vitamins, pigments, phenols and polyphenols. Synthetic antioxidants, such as butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA), are well known for their ability to stop the chain reactions of lipid peroxidation.

Unfortunately, it has been shown that they are able to enhance carcinogenesis, cause pulmonary damage in mice, liver necrosis, haemorrhagic death and neoplasia in rats¹⁷. In this respect, flavonoids and other polyphenols are paid the greatest attention due to their non-toxicity and potentiality for implementation in human diet. Therefore, the aim of this investigation was to analyze the composition of the essential leaf oil from *Cinnamomum zeylanicum* Blume from Sri Lanka and to test the antioxidant properties of the samples and the main compound eugenol using different methods to get informations about the antioxidative activities of this phenolic monoterpene itself and the as dominating component of the essential oil of cinnamon leaves.

Experimental

Materials: The essential leaf oil of *Cinnamomum zeylanicum* Blume from Sri Lanka (800830) and eugenol are products from Kurt Kitzing Co., Wallerstein, Germany.

Chemical composition of essential oil

Gas chromatographic (GC) analysis: GC-FID analyses were carried out using a GC-14A with SPME sleeve adapted to injector, FID and C-R6A-Chromatopac integrator (Shimadzu, Japan), a GC-3700 with FID (Varian, Germany) and C-R1B-Chromatopac integrator (Shimadzu). The carrier gas was hydrogen; injector temperature, 250°C; detector temperature, 320°C. The temperature programme was: 40°C/5 min to 280°C/5 min, with a heating rate of 6°C/min. The columns were 30 m x 0.32 mm bonded FSOT-RSL-200 fused silica, with a film thickness of 0.25 µm (Biorad, Germany) and 30 m x 0.32 mm bonded Stabilwax, with a film thickness of 0.50 µm (Restek, USA). Quantification was achieved using peak area calculations, and compound identification was carried out partly using correlations between retention times^{1,10,22-24}.

Gas chromatographic-mass spectroscopic (GC-MS) analysis: For GC-MS measurements a GC-17A with QP5000 (Shimadzu), SPME sleeve adapted to injector and Compaq-ProLinea data system (class5k-software), a GC-HP5890 with HP5970-MSD (Hewlett-Packard, USA) and ChemStation software on a Pentium PC (Böhm, Austria), a GCQ (Finnigan-Spectronex, Germany-USA) and Gateway-2000-PS75 data system (Siemens-Nixdorf, Germany, GCQ-software) were used. The carrier gas was helium; injector temperature, 250°C; interface-heating at 300°C, ion-source-heating at 200°C, EI-mode was 70 eV, and the scan-range was 41-450 amu. For other parameters, see description of GC/FID, above. Mass spectra correlations were done using Wiley, NBS, NIST and our own library as well as published data^{1,22,23}.

Antioxidant activity

Scavenging effect on 2,2-diphenyl-1-picryl hydrazyl radical (DPPH): The radical scavenging ability was determined according to the method of Mensor *et al.*³⁰. One ml of 0.3 mM alcoholic solution of DPPH was added to 2.5 ml of the samples with different concentration of cinnamon oil and eugenol. The samples were kept at room temperature in the dark and after 30 min the optic density was measured at 518 nm. The optic density of the samples, the control and the blank was measured in comparison with ethanol. One synthetic antioxidant, butylhydroxytoluene (BHT) and butylated hydroxyanisole (BHA) was used as positive control.

Detection of hydroxyl radicals by deoxyribose assay: The assay was performed as described by Halliwell *et al.*¹⁵, with minor changes. All solutions were freshly prepared. 1.0 ml of the reaction mixture contained 28 mM 2-deoxy-D-ribose (dissolved in KH₂PO₄/K₂HPO₄ buffer 10 mM, pH 7.4), 500 µl solution of various concentrations of the cinnamon oil or eugenol, 200 µM FeCl₃ and 1.04 mM EDTA (1:1 v/v), 10 mM H₂O₂ and 1.0 mM ascorbic acid. After an incubation period of 1 h at 37°C the extent of deoxyribose degradation was determined by the thiobarbituric acid (TBA) reaction. 1.0 ml of TBA (1 % in 50 mM NaOH) and 1.0 ml of trichloroacetic acid (TCA) were added to the reaction mixture and the tubes were heated at 100 °C for 20 min. After cooling, the absorbance was read at 532 nm against a blank (containing only buffer and deoxyribose). The percentage inhibi-

tion was calculated by the formula: $I(\%) = 100 - (\text{Abs}_{\text{sample}} / \text{Abs}_{\text{control}}) \times 100$.

The IC₅₀ value represented the concentration of the compounds that caused 50 % inhibition of radical formation. Quercetin was used as a positive control.

Evaluation of antioxidant activity in linoleic acid model system: Linoleic acid emulsions were prepared by mixing 0.285 g of linoleic acid, 0.289 g of Tween 20 as emulsifier and 50 ml phosphate buffer (pH 7.2). The mixture was homogenized for 5 min according to *et al.*⁴¹. The antioxidant was added at the final concentrations of 0, 0.0025, 0.005, and 0.01 % wt/vol of oil, BHT 0.01 % was used as control. The mixture was incubated in an oven at 37°C for 10 d. The progress of oxidation was monitored by measuring the conjugated diene formation (CD) and thiobarbituric acid reactive substances (TBARS).

The antioxidant activity at the end of assay time was expressed for each indicator as reduction percent of peroxidation (RP %) and for a control containing no antioxidant being 0%. $RP \% = [(\text{peroxidation indicator value without antioxidant}) - (\text{peroxidation indicator value with antioxidant}) / \text{peroxidation indicator value without antioxidant}] \times 100$. A higher percentage indicates a higher antioxidant activity.

Determination of conjugated dienes formation: Aliquots of 0.02 ml were taken at different intervals during incubation. After incubation, 2 ml of methanol in deionised water (60 %) were added, and the absorbance of the mixture was measured at 233 nm. The conjugated diene concentration was expressed in ml/mg in each sample. The results were calculated as $CD = B \times \text{vol/wt}$; where B is the absorbance reading, vol denotes the volume (ml) of the sample and wt is the mass (mg) of emulsion measured⁴².

Determination of thiobarbituric acid reactive substances: A modified thiobarbituric acid reactive substances (TBARS) method was used to measure the antioxidant activity of oil in terms of inhibition on lipid peroxidation. 0.1 ml of sample was taken every day, from the emulsion, the following were sequentially added: the TBA-TCA solution (20 mM TBA in 15 % TCA). The mixture was heated in a 100°C water bath for 15 min and cooled at room temperature. After 2 ml of chloroform were added, the mixture was mixed and centrifuged at 2000 rpm for 15 min. The chloroform layer was separated and the absorbance of the supernatant was measured at 532 nm against a blank containing TBA-TCA solution. Malonic aldehyde standard curves were prepared by 1,1,3,3-tetramethoxypropane and TBARS were expressed as mg of malonic aldehyde/kg dry matter.

The data obtained at each point for all experiments were the average of three measurements.

Statistical analysis: All experimental data (in triple repetition) were included in an approximation model through polynomial dependences from fourth order. For all cases the plural correlation coefficient R² was determined. The level of the concentration which corresponds to 50 % of inhibition was calculated according to this approximated dependence for which R² is maximum. The mathematical analysis of the data is carried out with specialized MatLab (5.3/6.0) software.

Results and Discussion

Chemical composition of the essential leaf oil of *Cinnamomum zeylanicum*

Blume from Sri Lanka: The essential oil of cinnamon leaves from Sri Lanka, possessing a pleasant cinnamon-leaf, eugenol- and clove-like as well as warm-floral scent, was analyzed by GC and GC-MS furnishing the following composition: Main compounds (concentrations higher than 2.0%, calculated as relative %-peak area of GC-FID analysis using an apolar column and mean-value of 3 measurements) are eugenol (74.9%), β -caryophyllene (4.1%), benzyl benzoate (3.0%), linalool (2.5%) and eugenyl acetate (2.1%). As further aroma compounds in medium concentrations (1.0-2.0%) cinnamyl acetate (1.8%), safrole (1.3%), α -pinene (1.2%) and cinnamaldehyde (1.1%) were identified. More than 20 lower concentrated volatiles are also aroma-active constituents of the *C. zeylanicum* sample (Table 1).

DPPH radical-scavenging activity: DPPH is a free radical compound that has been widely used to test the free radical-scavenging ability of various samples^{16,18,36}. Figure 1 depicts the scavenging activity of cinnamon oil, eugenol, BHT and BHA on DPPH radicals at various concentrations. Cinnamon essential oil demonstrated the highest inhibitory activity compared to other antioxidants studied, reaching as high as 94.42 % at 8.0 $\mu\text{g/ml}$, while for eugenol, BHT and BHA a concentration of at 20 $\mu\text{g/ml}$ was needed to achieve DPPH radicals' inhibition of 88.67, 82.40 and 89.98%, respectively. The concentration of cinnamon oil resulting in a 50 % inhibition of the free radical - IC_{50} , was 0.245 $\mu\text{g/ml}$, while for eugenol the corresponding concentration was 1.258 $\mu\text{g/ml}$. IC_{50} values were with statistical significance $p \leq 0.01$ and with high regression coefficients - $R^2 = 0.993$ (for cinnamon oil) and $R^2 = 0.999$ (for eugenol). The standards BHT and BHA showed IC_{50} of 4.407 $\mu\text{g/ml}$ ($R^2 = 0.998$) and 1.12 $\mu\text{g/ml}$ ($R^2 = 0.996$), respectively.

Hydroxyl radical-scavenging activity: Among all oxygen radicals the hydroxyl radical ($\text{OH}\cdot$) is the most reactive and damages diverse biomolecules. Hydroxyl radicals were generated in a reaction mixture containing ascorbate, hydrogen peroxide and iron III-EDTA at pH 7.4 and measured by their ability to degrade the sugar deoxyribose^{3,15}. Cinnamon oil showed high hydroxyl radical-scavenging activity that intensified with the increase of concentration, reaching 90.0% at 0.1 $\mu\text{g/ml}$, while at an equal concentration the dominant component of cinnamon oil – eugenol, caused 71.30 % inhibition of the hydroxyl radical (Figure 2). Substantially weaker was the antioxidant activity of quercetin – 77.8% at 20 $\mu\text{g/ml}$ concentration. The three assayed antioxidants were arranged by their antioxidant effect (expressed as IC_{50}) in descending order, as follows: cinnamon oil - 0.046 $\mu\text{g/ml}$ ($R^2 = 0.999$), eugenol - 0.055 $\mu\text{g/ml}$ ($R^2 = 0.989$) and quercetin - 4.61 $\mu\text{g/ml}$ ($R^2 = 0.834$). The same analytical method could also be applied for studying the inhibitory power of cinnamon oil against the metal ion-dependant generation of $\text{OH}\cdot$, and not only for assaying its ability to scavenge already formed radicals.

Heavy metals, particularly those possessing two or more valency states with a suitable oxidation-reduction potential between them (e.g. Fe), increase the maximum rate of lipid oxidation. Metals act as pro-oxidants by electron transfer, liberating radicals from fatty

acids or hydroperoxides. Chelation of metal ions by food components, particularly antioxidants, reduces the pro-oxidative effect of these ions and raises considerably the energy of activation of the initiation reactions. The efficiency of cinnamon oil (in comparison with quercetin and eugenol) in reducing the pro-oxidant effect of Fe-ions by chelation was tested.

When FeCl_3 was added to the reaction mixture the Fe^{3+} -ions formed a complex with deoxyribose and they may be subsequently reduced by ascorbate to Fe^{2+} , which further reacts with H_2O_2 . In a result, the $\text{OH}\cdot$ were generated and degradation of dextrose occurred. Only molecules that are able to chelate Fe ions and make them inactive may inhibit the degradation of deoxyribose. Results depicted on Figure 3 (without EDTA) show that cinnamon oil, eugenol and quercetin manifest chelative properties, most strongly expressed in the case of cinnamon oil.

Like most radicals, $\text{OH}\cdot$ can be neutralized by a hydrogen atom. The capture of $\text{OH}\cdot$ by cinnamon oil is attributed to the hydrogen-donating ability of the phenol eugenol, which is found in high concentrations (75.00 %) in cinnamon essential oil.

Evaluation of antioxidant activity in linoleic acid model system: For the purpose of evaluating the antioxidant activity of cinnamon oil an emulsion of linoleic acid was used as a model system. Cinnamon essential oil was tested for its ability to inhibit the generation of hydrogenperoxides at early stages of the oxidation of linoleic acid, as well as for its inhibitory potential after the emergence of secondary oxidized products, like aldehydes, ketones or hydrocarbons. The analyses of those properties were carried out using two indicators, corresponding to the different degrees of lipid peroxidation - conjugated diene formation and TBARS.

Determination of conjugated diene formation: It was determined that the peak in conjugated diene formation was attained on the 5th day following the incubation of linoleic acid (Fig. 4A). At the higher concentration applied - 0.01%, the inhibitory effect of cinnamon oil was comparable to the action of the synthetic antioxidant BHT, namely 56.52 % for cinnamon oil and 58.69 % for BHT, measured on the 9th day of the experiment. Cinnamon oil performed weaker inhibitory effect on lipid peroxidation at 0.005% concentration - 47.82% on the 9th day of storage.

Determination of thiobarbituric acid reactive substances: With the second indicator used - TBARS, peak accumulation of malonic aldehyde was observed, as above, on the 5th day of linoleic acid storage (Fig. 4B), suggesting that the process in concern ran almost parallel to the formation of conjugated dienes. The antioxidant action of 0.01% cinnamon oil was absolutely similar to that of the standard BHT at concentration of 0.01% - the inhibition of lipid peroxidation reached 75.88% in the case of cinnamon oil and 76.47% for BHT. The lower concentration of cinnamon oil under study - 0.005%, caused insignificantly weaker inhibition of the process, reaching 71.47% on the 8th day of the experiment.

Conclusion: The essential leaf oil of *Cinnamomum zeylanicum* Blume from Sri Lanka is rich in eugenol, benzyl benzoate, linalool and eugenyl acetate as well as further

aroma-active volatiles, responsible for the pleasant cinnamom and clove-like (strong eugenol-note) and warm-floral scent of this sample^{2,7,12,13,37}.

The scavenging of DPPH and OH• radicals by cinnamon oil is attributed to the hydrogen-donating capacity of the phenolic component eugenol, present in significant concentrations in the essential oil. Cinnamon oil manifested higher antioxidant activity towards the DPPH radical than eugenol, BHT and BHA, confirmed by the lowest IC₅₀ value. The inhibitory potential of cinnamon oil against OH• was displayed once again at a lower concentration of IC₅₀ than that of eugenol and quercetin. Cinnamon essential oil demonstrated high chelative activity with respect to the Fe³⁺, resulting in a prevention of hydroxyl radicals' initiation.

Cinnamon oil inhibited effectively the conjugated diene formation and the generation of secondary products from lipid peroxidation at concentrations equivalent to those of the standard BHT. Even more inhibited was the stage of the generation of secondary products of lipid autoxidation, i.e. cinnamon oil could also be applied as an antioxidant at a more advanced stage of lipid oxidation.

The use of some methods to determine the antioxidant properties of medicinal plant essential oils confirms the findings that antioxidant capacity detected by only a single method should be interpreted with some caution as published elsewhere²⁸.

In addition, our study demonstrates that cinnamon essential oil possesses considerable antioxidant capacity and could readily be implemented as a natural preservative, thus reducing or avoiding losses due to oxidative processes.

References

1. **Adams, R.P. (2001).** Identification of Essential Oil Components by Gas Chromatography/Quadruple Mass Spectroscopy. Allured Publishing Co., Carol Stream.
2. **Arctander, St. (1969).** Perfume and Flavor Chemicals (Aroma Chemicals). Arctander Publishing, Montclair.
3. **Aruoma, O., Laughton, M., Halliwell, B. (1989).** Carnosin, homocarnosin and anserine: could they act as antioxidants in vivo. *Biochem. J.* 264: 863-869.
4. **BACIS – Boelens Aroma Chemical Information Service. (2000).** Spices and Condiments, Cinnamom. VCF Volatile Compounds in Food Database. TNO Nutrition and Food Research, Zeist.
5. **Bahadori, B., Yazdani-Biuki, B. (2003).** Antioxidantien - Therapierichtlinien für die Praxis. LKH-Mürzzuschlag – Medizinische Universität Graz.
6. **Baratta, M., Dorman, H., Deans, S., Figueiredo, A., Barroso, J., Ruberto, G. (1998).** Antimicrobial and antioxidant properties of some commercial essential oils. *Flavour Fragr. J.* 13: 235-244.
7. **Bauer, K., Garbe, D., Surburg, H. (1997).** Common Fragrance and Flavor Materials, 3rd Ed., Wiley-VCH, Weinheim.
8. **Chericoni, S., Prieto, J.M., Iacopini, P., Cioni, P., Morelli, I. (2005).** In Vitro Activity of the Essential Oil of *Cinnamomum zeylanicum* and Eugenol in Peroxynitrite-Induced Oxidation Processes. *J. Agric. Food Chem.* 53: 4762-4765.
9. **Dang, M., Takacsova, M., Nguyen, D., Kristianova, K. (2005).** Antioxidant

- activity of essential oils from various spices, *Nahrung/Food*, 45: 64 – 66.
10. **Davies, N.W. (1990).** Gas chromatographic retention indices of monoterpenes and sesquiterpenes on methyl silicone and Carbowax 20M phases. *J. Chromatogr.* 503: 1-24.
 11. **Escobar, J.A., Rubio, M.A., Lissi A. (1995).** SOD and catalase inactivation by singlet oxygen and peroxy radicals. *Free Radic. Biol. Med.* 20: 285-290.
 12. **Fazzalari, F.A. (1978).** Compilation of Odor and Taste Threshold Values Data. American Society for Testing and Materials, Philadelphia.
 13. **Furia, T.E., Bellanca, N. (1975).** Fenaroli's Handbook of Flavor Ingredients, 2nd Ed., CRC Press, Cleveland.
 14. **Halliwell, B., Gutteridge, J.M.C. (1984).** Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem J.* 219: 1-14.
 15. **Halliwell, B., Gutteridge, J., Aruoma, O. (1987).** The deoxyribose method: a simple 'test tube' assay for determination of rate constants for reaction of hydroxyl radicals. *Anal. Biochem.* 165: 215-219.
 16. **Hatano, T., Takagi, M., Ito, H., Yoshida, T. (1997).** Phenolic constituents of liquorice. VII. A new calcone with a potent radical scavenging activity and accompanying phenols. *Chem. Pharm. Bull.* 45: 1485-1492.
 17. **Ito, N., Hirose, M., Fukushima, S., Tsuda, H., Shirai T., Tatematsu, M. (1985).** Studies of antioxidants: Their carcinogenic and modifying effect on chemical carcinogenesis. *Food Chem. Toxicol.* 24: 1071-1077.
 18. **Jadhav, H.R., Bhutani K.K. (2002).** Antioxidant properties of Indian medicinal plants. *Phytoter. Res.* 16: 771-773.
 19. **Jayaprakasha, G.K., Rao, L.J., Sakariah, K.K. (2002).** Chemical composition of volatile oil from *Cinnamomum zeylanicum* Buds. *Z. Naturforsch. C* 57: 990-993.
 20. **Jayaprakasha, G.K., Rao, L.J., Sakariah, K.K. (2003).** Volatile Constituents from *Cinnamomum zeylanicum* Fruit Stalks and Their Antioxidant Activities. *J. Agric. Food Chem.* 51: 4344-4348.
 21. **Jayaprakasha, G.K., Ohnishi-Kameyama, M., Ono, H., Yoshida, M., Rao, L.J. (2006).** Phenolic Constituents in the Fruits of *Cinnamomum zeylanicum* and Their Antioxidant Activity. *J. Agric. Food Chem.* 54: 1672-1679.
 22. **Jennings, W., Shibamoto, T. (1980).** Qualitative Analysis of Flavor and Fragrance Volatiles by Glass Capillary Gas Chromatography. Academic Press, New York.
 23. **Joulain, D., König, W.A. (1998).** The Atlas of Spectral Data of Sesquiterpene Hydrocarbons. E.B.-Verlag, Hamburg.
 24. **Kondjoyan, N., Berdaqué, J.L. (1996).** A Compilation of Relative Retention Indices for the Analysis of Aromatic Compounds. Edition du Laboratoire Flaveur. Saint Genes Champanelle.
 25. **Lee, K., Shibamoto, T. (2002).** Determination of antioxidant potential of volatile extracts isolated from various herb and spices, *J Agric. Food Chem.* 50: 4947-4952.
 26. **Lis-Balchin, M., Deans, S., Eaglesham, E. (1998).** Relationship between bioactivity and chemical composition of commercial essential oils. *Flavour Fragr. J.* 18: 98 - 104.

27. **Mancini-Filho, J., Van-Koij, A., Mancini, D.A.P., Cozzolino, F.F., Torres, R.P. (1998).** Antioxidant activity of cinnamom (*Cinnamomum zeylanicum*, Breyne) extracts. *Boll. Chim. Farm.* 137: 443-447.
28. **Mantle, D., Anderton, J.G., Falkous, G., Barnes, M., Jones, P., Perry, E.K. (1998).** Comparison of methods for determination of total antioxidative status: application to analysis of medicinal plant essential oils. *Biochem. Physiol.*, part B: *Biochem. Molecul. Biol.* 121B: 385-391.
29. **Mastura, M., Azah, M.A.N., Khozirah, S., Mawardi, R., Manaf, A.A. (1999).** Anticandidal and antidermatophytic activity of *Cinnamomum* species essential oils. *Cytobios* 98: 17-23.
30. **Mensor, L.L., Menezes, F.S., Leitao, G.G., Reis, A.S., Santos, T.S., Coube, C.S., Leitao, S.G. (2001).** Screening of Brazilian Plant Extracts for Antioxidant Activity by the Use of DPPH Free Radical Method. *Phytother. Res.* 15: 127-130.
31. **Murcia, M., Egea, I., Romajaro, F., Jimenes, P., Martinez-Tome, M. (2004).** Antioxidant evaluation in dessert spices compared with common food additives. Influence of irradiation procedure. *J. Agric. Food Chem.* 52: 1872-1881.
32. **Namiaki, M. (1990).** Antioxidants/antimutagens in food. *Food Sci. Nutr.* 29: 273-300.
33. **Ranjbar, A., Ghasmeinezhad, S., Zamani, H., Malekirad, A.A., Baiaty, A., Mohammadirad, A., Abdollahi, M. (2006).** Antioxidative stress potential of *Cinnamomum zeylanicum* in humans: a comparative cross-sectional clinical study. *Therapy* 3: 113-117.
34. **Seidemann, J. (2005).** Cinnamom – Lauraceae. *World Spice Plants*. Springer-Verlag, Heidelberg, 100-104.
35. **Shan, B., Cai, Y., Sun, M., Corke, H. (2005).** Antioxidant capacity of 26 extracts and characterization of their constituents. *J. Afric. Food Chem.* 53: 7749-7759.
36. **Shimoji, Y., Tamura, Y., Nakamura, Y., Nanda, K., Nishidai, S., Nishikawa, Y. (2002).** Isolation and identification of DPPH radical scavenging compounds in kurosu (Japanese unpolished rice vinegar). *J. Agric. Food Chem.* 50: 6501-6503.
37. **Sigma-Aldrich (2003).** *Flavors & Fragrances, The Essence of Excellence*. Sigma-Aldrich Co., Milwaukee.
38. **Teissedre, P.L., Waterhouse, A.L. (2000).** Inhibition of Oxidation of Human Low-Density Lipoproteins by Phenolic Substances in Different Essential Oils Varieties. *J. Agric. Food Chem.* 48: 3801-3805.
39. **Tomaino, A., Cimino, F., Zimbalatti, V., Venuti, V., Sulfaro, V., De Pusquale, A., Saiha, A. (2005).** Influence of heating on antioxidant activity and the chemical composition of some spice essential oils. *Food Chem.* 89: 549 – 554.
40. **Wichtl, M. (1989).** *Zimtrinde*. Teedrogen, 2nd. Ed., Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, 535-537.
41. **Yen, G., Chang, Y., Su, S. (2003).** Antioxidant activity and active compounds of rice koji fermented with *Aspergillus candidus*. *Food Chem.* 83: 49-54.
42. **Zainol, M., Abd-Hamid, A., Yusof, S., Muse, R. (2003).** Antioxidative activity and total phenolic compounds of leaf, root and petiole of four accessions of *Centella asiatica* (L.) Urban. *Food Chem.* 81: 575-581.

Table 1. Chemical composition of the essential leaf oil of *Cinnamomum zeylanicum* Blume from Sri Lanka.

Compound#	%+	RI*
<i>cis</i> -3-Hexenol	0.1	860
Styrene	0.1	897
α -Thujene	0.2	930
α -Pinene	1.2	939
Camphene	0.3	953
Benzaldehyde	0.1	961
β -Pinene	0.3	981
Myrcene	0.1	990
α -Phellandrene	0.9	1002
<i>cis</i> -3-Hexenyl acetate	0.1	1005
α -Terpinene	0.1	1015
<i>p</i> -Cymene	0.8	1024
Limonene	0.5	1026
β -Phellandrene	0.2	1028
δ -3-Carene	0.6	1030
1,8-Cineole	0.6	1032
Benzyl alcohol	0.2	1034
<i>trans</i> - β -Ocimene	0.1	1049
γ -Terpinene	0.1	1059
Terpinolene	0.2	1087
Linalool	2.5	1093
Phenylethyl alcohol	0.1	1105
Benzyl acetate	0.1	1160
α -Terpineol	0.3	1188
Cinnamaldehyde	1.1	1259
Safrole	1.3	1285
α -Cubebene	0.9	1350
Eugenol	74.9	1357
Isoeugenol	0.1	1405
β -Caryophyllene	4.1	1417
Cinnamyl acetate	1.8	1444
α -Humulene	0.6	1453
α -(<i>E,E</i>)-Farnesene	1.1	1508
Eugenyl acetate	2.1	1522
<i>trans</i> -Nerolidol	0.1	1561
<i>cis</i> -3-Hexenyl benzoate	0.1	1565
Caryophyllene oxide	0.5	1581
Benzyl benzoate	3.0	1758

#in order of their retention-times, +relative %-peak area of GC-FID analyses using an apolar column, *retention indices using an apolar OV-5-type column




Figure 1. Scavenging effects on DPPH radical of cinnamon, eugenol, BHT and BHA




Figure 2. Action of ocinnamon oil, eugenol and quercetin on hydroxyl radical – dependent degradation of deoxyribose.





Figure 3. Metal chelating activity of cinnamon oil, eugenol and quercetin on deoxyribose degradation by OH•






Figure 4. Effect of cinnamon oil on (A) conjugated dienes and (B) TBARS in a linoleic acid/water emulsion system